

# On $n$ -ary Lie algebras of type $(r, l)^1$

E.G. Vishnyakova

## Abstract

These notes are devoted to the multiple generalization of a Lie algebra introduced by A.M. Vinogradov and M.M. Vinogradov. We compare definitions of such algebras in the usual and invariant case. Furthermore, we show that there are no simple  $n$ -ary Lie algebras of type  $(n-1, l)$  for  $l > 0$ .

## 1 Introduction

This paper is devoted to the study of the multiple generalization of a Lie algebra introduced in [11]. More precisely in [11] A.M. Vinogradov and M.M. Vinogradov proposed a two-parameter family of  $n$ -ary algebras that generalized *Filippov  $n$ -algebras* and *Lie  $n$ -algebras*. V.T. Filippov [3] considered alternating  $n$ -ary algebras  $V$  satisfying the following Jacobi identity:

$$\{a_1, \dots, a_{n-1}, \{b_1, \dots, b_n\}\} = \sum_{i=1}^n \{b_1, \dots, b_{i-1} \{a_1, \dots, a_{n-1}, b_i\}, \dots, b_n\}, \quad (1)$$

where  $a_i, b_j \in V$ . In other words, the linear maps  $\{a_1, \dots, a_{n-1}, -\} : V \rightarrow V$  are derivations of the  $n$ -ary bracket  $\{b_1, \dots, b_n\}$ . Another natural  $n$ -ary generalization of the standard Jacobi identity has the following form:

$$\sum_{(I,J)} (-1)^{(I,J)} \{\{a_{i_1}, \dots, a_{i_n}\}, a_{j_1}, \dots, a_{j_{n-1}}\} = 0, \quad (2)$$

where the sum is taken over all ordered unshuffle multi-indexes  $I = (i_1, \dots, i_n)$  and  $J = (j_1, \dots, j_{n-1})$  such that  $(I, J)$  is a permutation of the multi-index  $(1, \dots, 2n-1)$ . We call alternating  $n$ -ary algebras satisfying Jacobi identity (2) *Lie  $n$ -algebras*.

The  $n$ -ary algebras of type (1) appear naturally in Nambu mechanics [9] in the context of Nambu-Poisson manifolds, in supersymmetric gravity theory and in supersymmetric gauge theories, the Bagger-Lambert-Gustavsson Theory, see [1] for details. The  $n$ -ary algebras of type (2) were considered for instance by P. Michor and A.M. Vinogradov in [8] and by P. Hanlon and M.L. Wachs [4]. The homotopy case was studied in [10] in the context of the Schlesinger-Stasheff homotopy algebras and  $L_\infty$ -algebras. Such algebras are related to the Batalin-Fradkin-Vilkovisky theory and to string field theory [6].

---

<sup>1</sup> Supported by AFR-grant, University of Luxembourg.

The paper is structured as follows. In Section 2 we remind the definition of the Nijenhuis-Richardson bracket on  $\bigwedge V^* \otimes V$ , where  $V$  is a finite dimensional vector space. In Section 3 we give a definition Vinogradovs' algebras. In Section 4 we compare definitions of usual Vinogradovs' algebras and Vinogradovs' algebras with a symmetric non-degenerate invariant form. In Section 4 we show that there are no simple  $n$ -ary Lie algebras of type  $(n-1, l)$ ,  $l > 0$ . The case of  $n$ -ary Lie algebras of type  $(n-1, 0)$  was studied in [7].

## 2 Nijenhuis-Richardson bracket

In this section we follow [11]. Let  $V$  be a finite dimensional vector space over  $\mathbb{R}$  or  $\mathbb{C}$ . We put  $I^n := (1, \dots, n)$  and we denote by  $I$  and  $J$  the ordered multi-indexes  $I = (i_1, \dots, i_l)$  and  $J = (j_1, \dots, j_k)$  such that  $i_1 < \dots < i_l$  and  $j_1 < \dots < j_k$ . Let  $|I| := l$  and  $|J| := k$ . Assume that  $i_p \neq j_q$  for all  $p, q$ . Then we have the non-ordered multi-index

$$(I, J) := (i_1, \dots, i_l, j_1, \dots, j_k).$$

We denote by  $I + J$  the ordering of the multi-index  $(I, J)$  and by  $(-1)^{(I, J)}$  the parity of the permutation that maps  $(I, J)$  into  $I + J$ . We set

$$a_{I^n} := (a_1 \dots, a_n) \quad a_I := (a_{i_1} \dots, a_{i_l}), \quad a_J := (a_{j_1} \dots, a_{j_k}).$$

Let us take  $L \in \bigwedge^l V^* \otimes V$  and  $K \in \bigwedge^k V^* \otimes V$ . We define:

$$L[K](a_{I^{l+k-1}}) = \sum_{I+J=I^{l+k-1}} (-1)^{(I, J)} L(K(a_I), a_J). \quad (3)$$

Here we assume that  $|I| = k$ ,  $|J| = l - 1$ .

**Definition 1** *The Nijenhuis-Richardson bracket*

$$[L, K]^{NR} \in \bigwedge^{l+k-1} V^* \otimes V,$$

of  $L \in \bigwedge^l V^* \otimes V$  and  $K \in \bigwedge^k V^* \otimes V$  is defined by the following formula:

$$[L, K]^{NR} := (-1)^{(l-1)(k-1)} L[K] - K[L].$$

For  $L \in \bigwedge^l V^* \otimes V$ , we denote by  $L_{a_1, \dots, a_p}$  the following  $(l-p)$ -linear map:

$$L_{a_1, \dots, a_p}(b_1, \dots, b_{l-p}) := L(a_1, \dots, a_p, b_1, \dots, b_{l-p}),$$

where  $a_i, b_j \in V$ .

### 3 Vinogradovs' algebras

In this subsection we recall the definition of  $n$ -ary algebras introduced by A.M. Vinogradov and M.M. Vinogradov in [11]. The graded versions of such algebras are studied in [12]. These  $n$ -ary algebras are a generalization of  $n$ -ary Lie algebras of type (1) and of type (2) for even  $n$ . Let  $V$  be a finite dimensional complex or real vector space.

**Definition 2** *An  $n$ -ary algebra structure  $\mu$  on  $V$  is an  $n$ -linear map:*

$$\mu : \underbrace{V \times \cdots \times V}_{n \text{ times}} \rightarrow V.$$

*An  $n$ -ary algebra is a pair  $(V, \mu)$ .*

Sometimes we will use the bracket notation  $\{a_1, \dots, a_n\}$  instead of  $\mu(a_1, \dots, a_n)$ .

**Definition 3** *An  $n$ -ary algebra is called skew-symmetric if*

$$\{a_1, \dots, a_i, a_{i+1}, \dots, a_n\} = -\{a_1, \dots, a_{i+1}, a_i, \dots, a_n\} \quad (4)$$

*for any  $a_p \in V$ .*

**Definition 4** ([11]) *Let  $(V, \mu)$  be an  $n$ -ary skew-symmetric algebra. We say that  $(V, \mu)$  is an  $n$ -ary Lie algebra of type  $(r, l)$  if the following holds:*

$$[\mu_{a_1, \dots, a_r}, \mu_{b_1, \dots, b_l}]^{NR} = 0, \quad (5)$$

*where  $a_i, b_j \in V$  and  $0 \leq l \leq r < n$ .*

**Example 1.** Filippov  $n$ -ary algebras or  $n$ -ary algebras satisfying (1) are exactly  $n$ -ary Lie algebras of type  $(n-1, 0)$ . For even  $n$ , the  $n$ -ary Lie algebras satisfying (2) are exactly  $n$ -ary algebras of type  $(1, 0)$ . (See [11] for details.)

**Remark 5** *The theory of Filippov  $n$ -ary algebras is relatively well-developed. For instance, there is a classification of simple real and complex Filippov  $n$ -ary algebras and an analog of the Levi decomposition [7]. W.X. Ling in [7] proved that there exists only one simple finite-dimensional  $n$ -ary Filippov algebra over an algebraically closed field of characteristic 0 for any  $n > 2$ . The simple Filippov  $n$ -ary superalgebras in the finite and infinite dimensional case were studied in [2]. It was shown there that there are no simple linearly compact  $n$ -ary Filippov superalgebras which are not  $n$ -ary Filippov algebras, if  $n > 2$ . A classification of linearly compact  $n$ -ary Filippov algebras was given in [2].*

**Remark 6** *The  $n$ -ary algebras of type (2) were studied for instance in [8] and [4]. In [13] the  $n$ -ary algebras of type (2) endowed with a non-degenerate invariant form were considered.*

## 4 Nijenhuis-Richardson and Poisson brackets on $V$

Let  $V$  be a finite dimensional vector space that is supplied with a non-degenerate symmetric bilinear form  $(,)$ . Then  $\bigwedge V$  possesses a natural structure of a Poisson superalgebra defined by the following formulas:

$$\begin{aligned} [x, y] &:= (x, y), \quad x, y \in W; \\ [v, w_1 \cdot w_2] &:= [v, w_1] \cdot w_2 + (-1)^{vw_1} w_1 \cdot [v, w_2], \\ [v, w] &= -(-1)^{vw} [w, v], \end{aligned}$$

where  $v, w, w_i$  are homogeneous elements in  $\bigwedge W$ . (See for example [5] or [13] for details.) Let us take any element  $\tilde{\mu} \in \bigwedge^{n+1} V$ . Then we can define an  $n$ -ary algebra structure on  $V$  in the following way:

$$\{a_1, \dots, a_n\} := [a_1, [\dots, [a_n, \tilde{\mu}] \dots]], \quad a_i \in V. \quad (6)$$

**Definition 7** *A skew-symmetric  $n$ -ary algebra structure is called invariant with respect to the form  $(,)$  if the following holds:*

$$(b, \{a_1, \dots, a_n\}) = -(a_1, \{b, a_2, \dots, a_n\}) \quad (7)$$

for any  $b, a_i \in V$ .

The following observation was noticed in [13]:

**Proposition 8** *Assume that  $V$  is finite dimensional and  $(,)$  is non-degenerate. Any skew-symmetric invariant  $n$ -ary algebra structures can be obtained by construction (6). In other words, for any skew-symmetric invariant  $n$ -ary algebra structure  $\mu \in \bigwedge^n V^* \otimes V$ , there exists  $\tilde{\mu} \in \bigwedge^{n+1} V$  such that the map  $\mu$  coincides with the map defined by the formula (6).*

Assume that  $(V, \mu)$  is an  $n$ -ary Lie algebra of type  $(r, l)$  and the algebra structure  $\mu$  is invariant with respect to  $(,)$ . It follows from Proposition 8 that the algebra  $(V, \mu)$  has another  $n$ -ary algebra structure  $\tilde{\mu} \in \bigwedge^{n+1}(V)$ . In this section we study the relationship between these two  $n$ -ary algebra structures. Since  $\mu$  and  $\tilde{\mu}$  define the same algebra structure on  $V$ , we have:

$$\mu(a_1, \dots, a_l) = [a_1, \dots, [a_l, \tilde{\mu}] \dots].$$

Let us take two skew-symmetric invariant  $n$ -ary algebra structures  $L \in \bigwedge^l V^* \otimes V$  and  $K \in \bigwedge^k V^* \otimes V$ . It follows from (3) that

$$L[K](a_{I+l+k-1}) = \sum_{I+J=I+l+k-1} (-1)^{(I,J)} [[a_I, \tilde{K}], [a_J, \tilde{L}]], \quad (8)$$

where  $|I| = k$ ,  $|J| = l - 1$  and for simplicity we denote

$$[a_I, \tilde{L}] := [a_{i_1}, \dots [a_{i_l}, \tilde{L}] \dots].$$

**Proposition 9** *Let us take two skew-symmetric invariant  $n$ -ary algebra structures  $L \in \bigwedge^l V^* \otimes V$  and  $K \in \bigwedge^k V^* \otimes V$ . Then we have:*

$$[a_{I^{l+k-1}}, [\tilde{L}, \tilde{K}]] = -(-1)^{(k+1)(l+1)} [L, K]^{NR}(a_{I^{l+k-1}}).$$

**Proof.** We set

$$a_{I^p} := (a_1, \dots, a_p); \quad [a_{I^p}, \tilde{\mu}] := [a_1, \dots [a_p, \tilde{\mu}]].$$

Furthermore, we have:

$$\begin{aligned} [a_{I^{l+k-1}}, [\tilde{L}, \tilde{K}]] &= \sum_{I+J=I^{l+k-1}} (-1)^{(I,J)+(k+1)(l+1)} [[a_I, \tilde{L}], [a_J, \tilde{K}]] \\ &+ \sum_{I'+J'=I^{l+k-1}} (-1)^{(I',J')+(l+1)k} [[a_{I'}, \tilde{L}], [a_{J'}, \tilde{K}]] \\ &= \sum_{I+J=I^{l+k-1}} (-1)^{(I,J)+(k+1)(l+1)} [[a_I, \tilde{L}], [a_J, \tilde{K}]] \\ &+ \sum_{I'+J'=I^{l+k-1}} (-1)^{(J',I')} [[a_{I'}, \tilde{L}], [a_{J'}, \tilde{K}]] \\ &= \sum_{I+J=I^{l+k-1}} (-1)^{(I,J)+(k+1)(l+1)} [[a_I, \tilde{L}], [a_J, \tilde{K}]] \\ &+ \sum_{I'+J'=I^{l+k-1}} -(-1)^{(J',I')} [[a_{J'}, \tilde{K}], [a_{I'}, \tilde{L}]], \end{aligned}$$

where  $|I| = l$ ,  $|J| = k - 1$  and  $|I'| = l - 1$ ,  $|J'| = k$ . By equation (8), we get

$$\begin{aligned} [a_{I^{l+k-1}}, [\tilde{L}, \tilde{K}]] &= (-1)^{(k+1)(l+1)} K[L](a_{I^{l+k-1}}) - L[K](a_{I^{l+k-1}}) \\ &= -(-1)^{(k+1)(l+1)} [L, K]^{NR}(a_{I^{l+k-1}}). \end{aligned}$$

The proof is complete.  $\square$

Let  $L \in \bigwedge^t V^* \otimes V$  be a skew-symmetric invariant  $n$ -ary algebra structures. Clearly,

$$L_{a_1, \dots, a_p} = c[a_1, \dots, [a_p, \tilde{L}]],$$

where  $c \neq 0$ . Proposition 9 implies the following. If a skew-symmetric invariant algebra  $(V, \mu)$  is given by an identity of type

$$[\mu_{a_1, \dots, a_r}, \mu_{a_1, \dots, a_l}]^{NR} = 0,$$

then the same algebra can be given by the following relation:

$$[[a_{I^r}\tilde{\mu}], [a_{I^l}\tilde{\mu}]] = 0.$$

The result of our study is the following.

**Theorem 10** *Let  $V$  be a finite dimensional vector space with a non-degenerate symmetric bilinear form  $(,)$  and  $\mu \in \bigwedge^n V^* \otimes V$  be an invariant algebra structure on  $V$ . Then  $(V, \mu)$  is an  $n$ -ary Lie algebra of type  $(r, l)$  if and only if the following holds:*

$$[[a_{I^r}\tilde{\mu}], [a_{I^l}\tilde{\mu}]] = 0$$

for all  $a_i, b_j \in V$ .

## 5 Algebras of type $(n - 1, l)$

In this section we consider  $n$ -ary algebras of type  $(n - 1, l)$ , where  $0 < l \leq n - 1$ . More precisely, we study  $n$ -ary skew-symmetric algebras satisfying (5) for  $k = n - 1$  and  $l > 0$ .

**Lemma 1.** Jacobi identity (5) of type  $(n - 1, l)$ , where  $0 \leq l \leq n - 1$ , is equivalent to the following identity:

$$\{a_1, \dots, a_{n-1}, \{b_1, \dots, b_n\}\} = \sum_{i=1}^{n-l} \{b_1, \dots, \{a_1, \dots, a_{n-1}, b_i\}, \dots, b_n\}. \quad (9)$$

**Proof.**

We will need the following formula, see [11]:

$$[L, K]_a^{NR} = [L_a, K]^{NR} + (-1)^{l-1} [L, K_a]^{NR},$$

where  $L \in \bigwedge^l V^* \otimes V$  and  $K \in \bigwedge^k V^* \otimes V$ . Let  $(V, \mu)$  is an  $n$ -ary Lie algebra of type  $(n - 1, l)$ . Let us take  $a_i, b_j, c_t \in V$  and denote  $L := \mu_{a_1, \dots, a_n}$  and  $K := \mu_{b_1, \dots, b_l}$ . We have

$$[L, K]_{c_1, \dots, c_k}^{NR} = \sum_i (-1)^{k-i} [L_{c_i}, K_{c_1, \dots, \hat{c}_i, \dots, c_k}]^{NR} + [L, K_{c_1, \dots, c_i, \dots, c_k}]^{NR}$$

where  $k = n - l$ . Further,

$$\begin{aligned} [L, K_{c_1, \dots, c_i, \dots, c_k}]^{NR} &= \{a_1, \dots, a_{n-1}, \{b_1, \dots, b_l, c_1, \dots, c_k\}\}; \\ [L_{c_i}, K_{c_1, \dots, \hat{c}_i, \dots, c_k}]^{NR} &= -\{b_1, \dots, b_l, c_1, \dots, \hat{c}_i, \dots, c_k, \{a_1, \dots, a_{n-1}\}\}. \end{aligned}$$

It is clear that  $[L, K]^{NR} = 0$  if and only if  $[L, K]_{c_1, \dots, c_k}^{NR} = 0$ . The last equality is equivalent to:

$$\begin{aligned}
\{a_1, \dots, a_{n-1}, \{b_1, \dots, b_l, c_1, \dots, c_k\}\} &= \\
\sum_{i=1}^k (-1)^{k-i} \{b_1, \dots, b_l, c_1, \dots, \hat{c}_i, \dots, c_k, \{a_1, \dots, a_{n-1}\}\}, \\
\{a_1, \dots, a_{n-1}, \{b_1, \dots, b_l, c_1, \dots, c_k\}\} &= \\
\sum_{i=1}^k \{b_1, \dots, b_l, c_1, \dots, \{a_1, \dots, a_{n-1}\}, \dots, c_k\}, \\
\{a_1, \dots, a_{n-1}, \{c_1, \dots, c_k, b_1, \dots, b_l\}\} &= \\
\sum_{i=1}^k \{c_1, \dots, \{a_1, \dots, a_{n-1}\}, \dots, c_k, b_1, \dots, b_l\}.
\end{aligned}$$

The proof is complete.  $\square$

**Example 2.** For instance, 2-ary Lie algebras of type  $(1, 1)$  are exactly the associative skew-symmetric algebras. Using the fact that these algebras are skew-symmetric, it is easy to see that all such algebras have the following property:  $\{\{a, b\}, c\} = 0$ . Therefore, this class is not reach.

**Definition 11** A vector subspace  $W \subset V$  is called an ideal of a skew-symmetric  $n$ -ary algebra  $(V, \mu)$  if  $\mu(V, \dots, V, W) \subset W$ .

For example  $\{0\}$  and  $V$  are always ideals of  $(V, \mu)$ . Such ideals are called *trivial*.

**Definition 12** An  $n$ -ary Lie algebra of type  $(n-1, l)$  is called simple if it is not 1-dimensional and it does not have any non-trivial ideals.

**Theorem 13** Let  $V$  be a Lie algebra of type  $(n-1, l)$ , where  $0 < l \leq n-1$ , over  $\mathbb{C}$ . Assume that  $V$  does not possess any non-trivial ideals. Then  $\dim V = 1$ . In other words, there are no simple algebras of type  $(n-1, l)$ , where  $0 < l \leq n-1$ .

**Proof.** Assume that  $\dim V > 1$  and  $V$  does not possess any non trivial ideals. Let us interchange  $b_{n-l}$  and  $b_{n-l+1}$  and use (9). We get:

$$\begin{aligned}
&\{a_1, \dots, a_{n-1}, \{b_1, \dots, b_{n-l-1}, b_{n-l+1}, b_{n-l}, b_{n-l+2}, \dots, b_n\}\} \\
&= \sum_{i=1}^{n-l-1} \{b_1, \dots, \{a_1, \dots, a_{n-1}, b_i\}, \dots, b_n\} \\
&+ \{b_1, \dots, b_{n-l-1}, \{a_1, \dots, a_{n-1}, b_{n-l+1}\}, b_{n-l}, b_{n-l+2}, \dots, b_n\}.
\end{aligned}$$

From this equation and (9), we have:

$$\begin{aligned} & \{b_1, \dots, b_{n-l-1}, \{a_1, \dots, a_{n-1}, b_{n-l}\}, b_{n-l+1}, \dots, b_n\} \\ &= \{b_1, \dots, b_{n-l}, \{a_1, \dots, a_{n-1}, b_{n-l+1}\}, b_{n-l+2}, \dots, b_n\}. \end{aligned} \quad (10)$$

Denote by  $\mathcal{L}$  the vector subspace in  $\text{End}(V)$  generated by  $\mu_{a_1, \dots, a_{n-1}}$ , where  $a_i \in V$ . Let  $D = \mu_{a_1, \dots, a_{n-1}}$  for some  $a_i \in V$ . We can rewrite (10) in the form:

$$\begin{aligned} & \{b_1, \dots, b_{n-l-1}, D(b_{n-l}), b_{n-l+1}, \dots, b_n\} \\ &= \{b_1, \dots, b_{n-l}, D(b_{n-l+1}), b_{n-l+2}, \dots, b_n\}. \end{aligned} \quad (11)$$

The  $n$ -ary algebra  $(V, \mu)$  is skew-symmetric and the equation (11) holds for any  $b_i$ . Hence, we can assume that the following equation holds

$$\{b_1, \dots, D(b_i), b_{i+1}, \dots, b_n\} = \{b_1, \dots, b_i, D(b_{i+1}), \dots, b_n\}$$

for any  $i$ . Applying this formula several times, we get:

$$\{b_1, \dots, D(b_i), b_{i+1}, \dots, b_n\} = \{b_1, \dots, b_{i+j-1}, D(b_{i+j}), \dots, b_n\}$$

for all  $i, j$ . Furthermore, let us take  $D_1, D_2 \in \mathcal{L}$ . First of all, we see:

$$\begin{aligned} \{b_1, \dots, D_1(D_2(b_i)), b_{i+1}, \dots, b_n\} &= \{b_1, \dots, D_2(b_i), \dots, D_1(b_{i+j}), \dots, b_n\} \\ &= \{b_1, \dots, b_i, \dots, D_2(D_1(b_{i+j})), \dots, b_n\}. \end{aligned}$$

We put  $B := D_1 \circ D_2 - D_2 \circ D_1$ . Therefore,

$$\{b_1, \dots, B(b_i), \dots, b_{i+j}, \dots, b_n\} = -\{b_1, \dots, b_i, \dots, B(b_{i+j}), \dots, b_n\}.$$

Furthermore,

$$\begin{aligned} \{B(b_1), b_2, b_3, \dots, b_{i+1}, \dots, b_n\} &= -\{b_1, B(b_2), b_3, \dots, b_{i+1}, \dots, b_n\} \\ &= \{b_1, b_2, B(b_3), \dots, b_{i+1}, \dots, b_n\}; \\ \{B(b_1), b_2, b_3, \dots, b_{i+1}, \dots, b_n\} &= -\{b_1, b_2, B(b_3), \dots, b_{i+1}, \dots, b_n\}. \end{aligned}$$

Therefore,

$$\{B(b_1), b_2, b_3, \dots, b_n\} = 0$$

for all  $b_i \in V$ . If  $B(b_1) \neq 0$ , we see that  $\langle B(b_1) \rangle$  is a non-trivial 1-dimensional ideal in  $V$ . Hence,  $B(b_1) = 0$  for all  $b_1 \in V$ . Therefore,  $D_1 \circ D_2 - D_2 \circ D_1 = 0$  and  $\mathcal{L}$  is a usual abelian Lie algebra. Since,  $V$  does not possess any non-trivial ideals, the  $\mathcal{L}$ -module  $V$  should be irreducible. By Schur's Lemma  $\mathcal{L} \subset \{a \text{ id}\} \subset \text{End } V$  and  $\dim V = 1$ .  $\square$

**Example 3.** The classification of simple complex and real Filippov  $n$ -ary algebras or Lie algebras of type  $(n-1, 0)$  was done in [7]: there is one series of complex Filippov  $n$ -ary algebras  $A_k$ , where  $k$  is a natural number, and several real forms for each  $A_k$ . All these algebras have invariant forms and in terms of Poisson bracket they are given by the top form of  $V$ .

## References

- [1] de Azcarraga J.A., Izquierdo J.M., *n-ary algebras: a review with applications*, Journal of Physics A Mathematical and Theoretical, 43, 293001, 2010.
- [2] Cantarini N., and Kac V.G., *Classification of Simple Linearly Compact n-Lie Superalgebras*, Communications in Mathematical Physics 298.3 (2010): 833-853.
- [3] Filippov V. T., *n-ary lie algebras*, Sibirskii Math. J. 24 (1985) 126-140.
- [4] Hanlon P., Wachs M. L., *On Lie k-algebras*, Adv. in Math. 113 (1995) 206-236.
- [5] Kostant, B., Sternberg S., *Symplectic reduction, BRS cohomology and infinite-dimensional Clifford algebras*, Annals of Physics, 176:49-113, 1987.
- [6] Lada T., Stasheff J., *Introduction to SH Lie algebras for physicists*, Int. J. Theor.Phys. 32 (1993) 1087-1104, [arXiv:hep-th/9299099v1](#).
- [7] Ling W., *On the structure of n-Lie algebras*. Dissertation, University-GHS-Siegen, Siegn, 1993.
- [8] Michor P., Vinogradov A.M, *n-ary Lie and Associative Algebras. Geometrical structures for physical theories*, II (Vietri, 1996). Rend. Sem. Mat. Univ. Politec. Torino 54 (1996), no. 4, 373-392.
- [9] Nambu, Y., *Generalized Hamiltonian dynamics*, Phys. Rev. (1973), 2405-2412.
- [10] Schlesinger M., Stasheff J. D., *The Lie algebra structure of tangent cohomology and deformation theory*, J. Pure Appl. Algebra 38 (1985) 313-322.
- [11] Vinogradov A. M. and Vinogradov M. M., *On multiple generalizations of Lie algebras and Poisson manifolds*, 1998.
- [12] Vinogradov A.M. and Vinogradov M.M., *Graded Multiple Analogs of Lie Algebras*, Acta Applicandae Mathematicae 72: 183197, 2002.
- [13] Vishnyakova E.G., *Symmetric n-ary superalgebras with skew-symmetric invariant forms*, J.Geom.Phys. 98 (2015) 340-354.

*Elizaveta Vishnyakova*

Max Planck Institute for Mathematics Bonn and

University of Luxembourg

*E-mail address:* VishnyakovaE@googlemail.com